

PEDIATRIC IMMUNIZATIONS – SHARED DECISION MAKING

IMMUNIZATIONS	Birth-1 m	2 m	4 m	6 m	12 m	15 m	18 m	4 y	11 y	By 12 y	16 y	by 17 y	18 y	Required Daycare?	Required School?	What is this for?	Common reaction(s)
Hepatitis B	X													Yes	No	>For a blood borne infection that causes liver illness and flu-like symptoms	none
Pediarix - three components: <ul style="list-style-type: none"> • DTaP (diphtheria-tetanus-pertussis) • Hepatitis B • IPV (inactive polio) 		X	X	X										Yes	Yes DTaP/IPV No Hep B	>For a deadly throat infection/infection related to dirty injury/whooping cough > For a blood borne infection that causes liver illness and flu-like symptoms >For an illness that causes permanent muscle damage	Fever, redness/fullness/bump at the site of injection, fussiness, sleepiness at the 2 and 4 month times
Hemophilus Influenza B (HIB)		X	X				X							Yes	No	>For a serious infection around the brain that infants can get	none
Pneumococcal (Prevnar)		X	X	X		X								Yes	No	>For childhood ear infections	Fever, redness/fullness/bump at the site of injection, fussiness
Rotavirus (Rotateq)		X	X	X										No	No	>For an intestinal infection that makes infants sick and dehydrated	Gas, mild diarrhea
MMR (measles-mumps-rubella)					X									Yes	Yes	>For three viral illnesses that can cause serious infections and can cause male sterility	Fever 7-10 days after injection, redness/fullness/bump at the site of injection, mild body rash 7-10 days after, painful injection
Varicella (Varivax)					X									Yes	Yes	> For childhood infection with chicken pox	Fever 7-10 days after injection, redness/fullness/bump at the site of injection, mild body rash 7-10 days after, painful injection
Hepatitis A					X		X							Yes	No	>For an infection of the liver with flu like symptoms that can be passed through tainted foods	none
DTaP (diphtheria-tetanus-pertussis)						X								Yes	Yes	>For a deadly throat infection/infection	Mild flu symptoms, pain/redness/ bump at



IMMUNIZATIONS	Birth-1 m	2 mo	4 mo	6 mo	12 mo	15 mo	18 mo	4 y	11 y	12 y	16 y	17 y	18 y	Required Daycare	Required School?	related to dirty injury/whooping cough What is this for?	the site of injection, sore arm Common reaction(s)
MMR-V (Proquad) <ul style="list-style-type: none"> MMR Varicella 								X						Yes	Yes	>For three viral illnesses that can cause serious infections and can cause male sterility > For childhood infection with chicken pox	Mild fever, 7-10 days after injection, redness/fullness/bump at the site of injection, mild body rash 7-10 days after, sore arm, painful injection
DTaP-IPV (Kinrix) <ul style="list-style-type: none"> DTaP (diphtheria-tetanus-pertussis) IPV (inactive polio) 								X						Yes	Yes	>For a deadly throat infection/infection related to dirty injury/whooping cough >For an illness that causes permanent muscle damage	Mild flu symptoms, pain/redness/ bump at the site of injection, sore arm
Tdap – tetanus booster									X				X	Yes	Yes	BOOSTER FOR DTaP	Sore arm, redness/fullness/bump at the site of injection
Meningitis (Menveo/MCV4)									X		X			Yes	Yes	>For a deadly infection of the brain common in the teenage and early adult years	Sore arm, mild redness at the site of injection
Human Papilloma (HPV)									X	X				No	No	>For cervical cancer and genital warts, sexually transmitted, best given prior to sexual activity	Sore arm, can faint after injection
Meningitis B (Bexero)											X	X		No	No	>For a deadly infection of the brain common in the teenage and early adult years	Sore arm, mild redness at the site of injection
TOTAL INJECTIONS FOR THIS VISIT	1	3	3	2	3	2	2	2	3	1	2	1	1				

****What to do to help your child with the common reactions to shots (usually for the first 24 hours):**

Medications: Tylenol as needed (for fussiness or fever) every 4 hours

Massage: Massage the area gently where the injections were given (a warm bath helps as well)

Move: Move the extremity gently several times a day

Call the office for any concerns: 784-5782

